arXiv:1208.1281v1 [hep-ph] 6 Aug 2012

Model-Independent Evolution of Transverse Momentum Dependent Distribution Functions (TMDs) at NNLL

Miguel G. Echevarría,[1, *] Ahmad Idilbi,[2, †] Andreas Schäfer,[2, ‡] and Ignazio Scimemi[1, §]
[1]*Departamento de Física Teórica II, Universidad Complutense de Madrid (UCM), 28040 Madrid, Spain*
[2]*Institüt für Theoretische Physik, Universität Regensburg, D-93040 Regensburg, Germany*

We present a new approach to treat the perturbative QCD evolution of TMDs by resummation. We obtain a universal evolution kernel for the eight quark distributions while large logarithms are resummed up to next-to-next-to leading logarithms (NNLL). Contrary to earlier works this resummation gives unambiguous predictions avoiding the need to introduce an ad-hoc cut-off parameters and thus highly reducing the model-dependence. Applications to single-spin asymmetries, as the Sivers function, and unpolarized transverse-momentum-dependent parton distribution functions are given. The results agree very well with phenomenology.

PACS numbers:

Introduction: To formulate a complete, consistent theory of TMDs is one of the great challenges in modern QCD. TMDs are needed for all processes for which intrinsic transverse parton momenta are relevant, which form a large group. For example spin-dependent transverse momentum asymmetries provide unique clues to clarify the internal spin, angular momentum and 3-dimensional structure of hadrons. While operating at different energies, experiments and facilities such as HERMES, COMPASS, JLab and BNL are pursuing intensive research programs to explore TMDs. On the theoretical side, Sivers [1] and Collins [2] asymmetries have been intensely studied (see [3] for a review of TMD functions in spin-physics), and have attracted much attention recently [4–10] in context with fundamental theoretical concepts. Basically, some of the observed spin-asymmetries are linked to the presence of gauge links in non-collinear non-local correlators needed to maintain gauge invariance.

A complete formalism for the treatment of TMDs is still being developed. In this paper we refer to two approaches, namely the one developed by three of us in [11] and the one developed mainly by Collins [12]. For certain choices of parameter(s) one can argue that the two are equivalent, despite the technical differences. Here we address a long-standing problem of model-dependence and show how it is resolved by a new resummation technique which relies, in part, on the extracted anomalous dimension of the unpolarized quark-TMD up to $\mathcal{O}(\alpha_s^3)$, as was explained in [11].

The study of the unpolarized TMDs was pioneered by Collins and Soper [13, 14]. Collins' approach to TMDs consists of properly defining those quantities in a way consistent with a generic factorization theorem, extracting their anomalous dimensions and their evolution properties. This approach relies mainly on taking some of the Wilson lines in the soft factor off-the-light-cone. When doing so, one introduces an auxiliary parameter ζ specifying the measure of "off-the-light-coness". A differential equation with respect to ζ, the Collins-Soper evolution equation is then derived and solved which resums large logarithms and determines the evolution of the non-perturbative TMDs with energy.

Though Collins' approach is theoretically robust, two main drawbacks of it affect its phenomenological applications. First is the difficulty to perform off-the-light-cone calculations (beyond $\mathcal{O}(\alpha_s)$) compared to the on-the-light-cone ones. This, in practice, limits the accuracy to which large logarithms can be resummed. Second, and in order to avoid the Landau pole, an ad hoc non-perturbative model has to be introduced, which depends on an arbitrary parameter $b_{\max}$ that can only be extracted through fitting to experimental data. It turns out that the choice of $b_{\max}$ affects the evolution of TMDs *not* only at very low energies but at intermediate ones as well. This can be clearly noticed from the plots introduced below.

In this letter we present a new resummation technique based on the formalism developed in [11]. With what we call the "resummed D", explained below, we avoid the issue of Landau pole in a way where there is no need to introduce any ad hoc (cut-off) parameter, thereby eliminating the model-dependence to a precision far exceeding all current practical purposes. The parameter-free prediction of our formalism agrees perfectly with the results obtained for the phenomenologically preferred value of b_{max} [15]. We see this as an important step towards a complete theory of TMDs.

Definition of Quark-TMDs: Extending the work done in [11], we define a quark-TMD of a polarized hadron, collinear with the $+z$ direction with momentum P and spin $\vec{S}$ as

$$F_{\alpha\beta}(x,\vec{k}_\perp) = \frac{1}{2}\int \frac{dr^- d^2\vec{r}_\perp}{(2\pi)^3} e^{-i(\frac{1}{2}r^- xP^+ - \vec{r}_\perp\cdot\vec{k}_\perp)} \times \Phi^q_{\alpha\beta}(0^+,r^-,\vec{r}_\perp)\sqrt{S(0^+,0^-,\vec{r}_\perp)}\,. \quad (1)$$

$\Phi^q_{\alpha\beta}$ stands for a purely collinear matrix element, i.e., a matrix element which has no overlap with the soft region [16], and it is given by the bilocal correlator

$$\Phi^q_{\alpha\beta} = \langle P\boldsymbol{S}|\ \left[\bar{\xi}_{n\alpha} W^T_n\right](0^+,y^-,\vec{y}_\perp)\ \left[W^{T\dagger}_n \xi_{n\beta}\right](0)\,|P\boldsymbol{S}\rangle\,. \quad (2)$$

The soft function S which encodes soft-gluon emission is given by

$$S = \langle 0|\,\mathrm{Tr}\,\left[S^{T\dagger}_n S^T_{\bar{n}}\right](0^+,0^-,\vec{y}_\perp)\left[S^{T\dagger}_{\bar{n}} S^T_n\right](0)\,|0\rangle\ . \quad (3)$$

We should mention that $F_{\alpha\beta}$ is free from rapidity divergences, thus it is independent of any parameter like ζ.

One can obtain the eight leading-twist quark-TMDs [17, 18], represented generically by F below, simply by tracing $F_{\alpha\beta}$ with the Dirac structures $\frac{\bar{n}\!\!\!/}{2}$, $\frac{\bar{n}\!\!\!/\gamma_5}{2}$ and $\frac{i\sigma^{j+}\gamma_5}{2}$ for unpolarized, linearly polarized and transversely polarized quarks inside a polarized hadron. The superscript T indicates transverse gauge-links $T_{n(\bar{n})}$, necessary to render the matrix elements gauge-invariant [19, 20]. The definitions of collinear $(W_{n(\bar{n})})$, soft $(S_{n(\bar{n})})$ and transverse $(T_{n(\bar{n})})$ Wilson lines for DY and DIS kinematics can be found in [11].

In [11] the anomalous dimension of the unpolarized TMD was given up to 3-loop order based on a factorization theorem for q_T-dependent observables in a Drell-Yan process. In impact parameter space such factorization theorem for the hadronic tensor can be schematically written as: $\tilde{W} = H\tilde{F}_{f/P}\tilde{F}_{\bar{f}/\bar{P}}$, where H is the hard coefficient which encodes the physics at the probing scale Q. This quantity is built, order by order in perturbation theory, by considering virtual Feynman diagrams only, i.e., no real gluon emission has to be considered. Moreover, the quantity H has to be free from infrared physics, no matter how the latter is regularized. This should be the case whether one works on or off-the-light-cone.

Since the factorization theorem given above holds, at leading twist, also for spin-dependent observables, one can apply the same arguments, based on renormalization group invariance, as for the unpolarized case, to get a relation between the anomalous dimensions of $\tilde{F}$ and H, i.e., $\gamma_F = -\frac{1}{2}\gamma_H$, where $\gamma_H = 2\Gamma_{\rm cusp}\ln(Q^2/\mu^2) + 2\gamma_V$ is known at 3-loops level [21, 22]. This crucial result can be automatically extended to all TMDs defined in Eq. (1), since the anomalous dimension is independent of spin structure.

Evolution Kernel: For spin-dependent TMDs the operator product expansion for collinear PDFs fails. One has to resort to non-perturbative models, fitted at low energies to experimental data, to make predictions for higher energy experimental probes. Obviously, knowing the evolution kernel to as high accuracy as possible is key.

Starting from Eq. (1) the evolution of a generic quark-TMD from an initial scale Q_i to a final scale Q_f is given by $\tilde{F}(x,b;Q_f) = \tilde{F}(x,b;Q_i)\,\tilde{R}(b;Q_i,Q_f)$ where the evolution kernel $\tilde{R}$ is [8, 11, 12]

$$\tilde{R}(b;Q_i,Q_f) = \exp\left\{\int_{Q_i}^{Q_f}\frac{d\bar{\mu}}{\bar{\mu}}\gamma_F\left(\alpha_s(\bar{\mu}),\ln\frac{Q_f^2}{\bar{\mu}^2}\right)\right\}\left(\frac{Q_f^2}{Q_i^2}\right)^{-D(b;Q_i)}\,. \quad (4)$$

The D term can be obtained by noticing that the renormalized F has to be well-defined when its partonic version is calculated pertubatively. This means that all divergences, other than the genuine long-distance ones has to cancel. This fundamental statement – that rapidity divergences cancel when the collinear and soft matrix elements are combined according to Eq. (1)– allows one to extract all the Q^2-dependence from the TMDs and exponentiate it with the function D [11] thereby summing large logarithms of the form $\ln(Q^2/q_T^2)$. Applying the renormalization group invariance to $\tilde{W}$ we get $dD/d\ln\mu = \Gamma_{\rm cusp}$ by which we extract, for all the quark spin-dependent TMDs the function D at NNLO from the known cusp anomalous dimension at three-loops [22]. In this way we do not have to solve the Collins-Soper equation.

The function D contains large logarithms of the form $L_\perp = \ln\frac{\mu^2 b^2}{4e^{-2\gamma_E}}$ that have to be resummed. Matching the expansion $D(b,\mu) = \sum_{n=1}^\infty d_n(L_\perp)\left(\frac{\alpha_s}{4\pi}\right)^n$ to the expansions $\Gamma_{\rm cusp}(\alpha_s) = \sum_{n=1}^\infty \Gamma_{n-1}\left(\frac{\alpha_s}{4\pi}\right)^n$ and $\beta(\alpha_s) = -2\alpha_s\sum_{n=1}^\infty \beta_{n-1}\left(\frac{\alpha_s}{4\pi}\right)^n$ one gets the following recursive differential equation

$$d'_n(L_\perp) = \frac{1}{2}\Gamma_{n-1} + \sum_{m=1}^{n-1} m\beta_{n-1-m}d_m(L_\perp)\,, \quad (5)$$

where $d'_n \equiv dd_n/dL_\perp$. Solving this equation we can express the coefficients d_n in the following way,

$$2d_n = \frac{\Gamma_0}{\beta_0}\frac{(\beta_0 L_\perp)^n}{n} + (\beta_0 L_\perp)^{n-1}\left(\frac{\Gamma_0\beta_1}{\beta_0^2}\left(-1+H^{(1)}_{n-1}\right)|_{n\geq 3} + \frac{\Gamma_1}{\beta_0}|_{n\geq 2}\right) + (\beta_0 L_\perp)^{n-2}\left((n-1)d_2^0|_{n\geq 2} + (n-1)\frac{\Gamma_2}{2\beta_0}|_{n\geq 3} + \frac{\beta_1\Gamma_1}{\beta_0^2}s_n|_{n\geq 4} + \frac{\beta_1^2\Gamma_0}{\beta_0^3}t_n|_{n\geq 5} + \frac{\beta_2\Gamma_0}{2\beta_0^2}(n-3)|_{n\geq 4}\right) + \ldots\,, \tag{6}$$

where

$$\begin{aligned} s_n &= (n-1)H^{(1)}_{n-2} + \frac{1}{2}(5-3n)\,,\\ t_n &= \frac{1}{2}\Big[(1-n)H^{(2)}_{n-1} + n + 1\\ &\quad + (n-1)(\psi(n)+\gamma_E-2)(\psi(n)+\gamma_E)\Big]\,. \end{aligned} \tag{7}$$

$H^{(r)}_n$ is the r-th order Harmonic Number function of n and $\psi(n)$ is the digamma function of n. We can perform the resummation of large logs in the D when $\alpha_s L_\perp \sim 1$ and obtain with $a = \alpha_s/(4\pi)$ and $X = a\beta_0 L_\perp$

$$\begin{aligned} 2D &= -\frac{\Gamma_0}{\beta_0}\ln(1-X) + a\left[-\frac{\beta_1\Gamma_0}{\beta_0^2}\frac{X+\ln(1-X)}{(1-X)} + \frac{\Gamma_1}{\beta_0}\frac{X}{1-X}\right] + a^2\Big[d_2^0\frac{1}{(1-X)^2} + \frac{\Gamma_2}{2\beta_0}\frac{X(2-X)}{(1-X)^2}\\ &\quad + \frac{\beta_1\Gamma_1}{\beta_0^2}\frac{(X(X-2)-2\log(1-X))}{2(1-X)^2} + \frac{\beta_2\Gamma_0}{2\beta_0^2}\frac{X^2}{(1-X)^2}\\ &\quad - \frac{\beta_1^2\Gamma_0}{\beta_0^3}\frac{-121X^6+188X^5-13X^4-30X^3}{24(X-1)^2X^2}\\ &\quad - \frac{\beta_1^2\Gamma_0}{\beta_0^3}\frac{-12X^2(1-\mathrm{Li}_2(X))-12X(X+1)\log(1-X)}{24(X-1)^2X^2}\\ &\quad + \frac{\beta_1^2\Gamma_0}{2\beta_0^3}\sum_{n=5}^{\infty}X^{n-2}(n-1)\left[H^{(1)}_{n-1}\right]^2\Big] + \ldots \end{aligned} \tag{8}$$

For the last term in this expression we have found an analytic form of the sum of the series by using the approximation $H^{(1)}_{n-1} = \ln(n) + \gamma_E + \frac{1}{2n} - \frac{1}{12n^2} + \frac{1}{120n^4} - \frac{1}{256n^6} + \ldots$ which is precise enough for our purposes.

This resummed series is valid for $|X| < 1$, however when analytically continued through Borel-summation its validity is extended to $X \to -\infty$, which corresponds to $b \to 0$. The maximum value of b where this series can be used, corresponding to $X = 1$, is $b_c = 2e^{-\gamma_E}/\mu \exp[2\pi/(\beta_0\alpha_s(\mu))]$, and takes 8.42 GeV^{-1} for $\mu = Q_i = \sqrt{2.4}$ GeV.

For all practical applications, the resummed D will determine the evolution kernel, $\tilde{R}$, for all polarized and unpolarized TMDs. Moreover we are able to resum large logarithms up to NNLL accuracy.

Next, we briefly present Collins approach and illustrate the differences with ours by considering unpolarized quark TMD and Sivers function.

Within Collins approach the evolution kernel is [5, 12]

$$\tilde{R}^{\mathrm{C}}(b;Q_i,\zeta_i,Q_f,\zeta_f) = \exp\left\{-g_K(b)\ln\frac{\sqrt{\zeta_f}}{\sqrt{\zeta_i}}\right\} \times\exp\left\{\int_{Q_i}^{Q_f}\frac{d\bar{\mu}}{\bar{\mu}}\gamma_F^{\mathrm{C}}\left(\alpha_s(\bar{\mu}),\ln\frac{\zeta_f}{\bar{\mu}^2}\right)\right\}\left(\frac{\sqrt{\zeta_f}}{\sqrt{\zeta_i}}\right)^{\tilde{K}\left(b^*,\sqrt{\zeta_i}\right)}\,, \tag{9}$$

where the Collins-Soper kernel $\tilde{K}$ is resummed through its RG-equation

$$\tilde{K}\left(b^*;\sqrt{\zeta_i}\right) = \tilde{K}(b^*;\mu_b) - \int_{\mu_b}^{\sqrt{\zeta_i}}\frac{d\bar{\mu}}{\bar{\mu}}\gamma_K\,. \tag{10}$$

Large $L_\perp$ logarithms are cancelled by choosing $\mu_b = 2e^{-\gamma_E}/b^*$ [23]. To avoid the Landau pole that appears when Fourier transforming back to momentum space, one writes $b^* = b/\sqrt{1+(b/b_{\max})^2}$ instead of b. $b_{\max}$ is an arbitrary cutoff that accounts for the separation between perturbative and non-perturbative regimes. Below we have implemented the BLNY model to compare our approach with Collins' with the following: $g_K(b) = \frac{g_2}{2}b^2$, and $g_2 = 0.68\ \mathrm{GeV}^2$ for $b_{\max} = 0.5\ \mathrm{GeV}^{-1}$ [24] and $g_2 = 0.184\ \mathrm{GeV}^2$ for $b_{\max} = 1.5\ \mathrm{GeV}^{-1}$ [15].

Within Collins' approach one usually identifies $\zeta_i = Q_i^2$ and $\zeta_f = Q_f^2$. With this choice one gets at $\mathcal{O}(\alpha_s)$: $\gamma_F^{\mathrm{C}} = \gamma_F$ and $\tilde{K} = -2D$, and those relations should hold to all orders in perturbation theory. However, γ_F^{C} beyond one-loop is not known and its determination would require an extremely difficult perturbative calculations.

Results and Conclusions: In Fig. 1 we show the evolution kernel from $Q_i = \sqrt{2.4}$ GeV to $Q_f = 5$ GeV with Collins' and our approaches. The impact of the differences on the evolution of two TMDs is illustrated in Fig. 2.



FIG. 1: *Evolution kernel from* $Q_i = \sqrt{2.4}$GeV *up to* $Q_f =$ 5GeV *and* $Q_f = 91.19$GeV *with different approaches.*



FIG. 2: *Up quark unpolarized TMD and Sivers function (Bochum and Torino fits shown with opposite sign) evolved from* $Q_i = \sqrt{2.4}$GeV *up to* $Q_f = 5$GeV *with different approaches to the evolution kernel.*

In order to perform the resummation of large logarithms consistently up to N^{i}LL order (or N^{i-1}LO in RG-improved perturbation theory) one needs the input shown in table I. In our approach one takes the resummed series in Eq. (8) up to the corresponding order i. In [4, 5, 8] γ_K was not implemented at 2-loop order, as it should be to get a complete NLL result.

Order	Accuracy $\sim \alpha_s^n L^k$	γ_V	Γ_{cusp}	D
N^{i}LL	$n+1-i \leq k \leq 2n$ (α_s^{i-1})	α_s^i	α_s^{i+1}	α_s^i

TABLE I: Approximation schemes for the resummed TMD, where $L = \ln(Q^2/\mu^2)$ and α_s^i indicates the order of the perturbative expansion.

The unpolarized quark-TMD at low energy is modeled as a Gaussian: $f_{up/P}(x;Q_i)\exp[-\sigma b_T^2]$ with $\sigma = 0.38/4\,\mathrm{GeV}^2$ [25], and we have taken the MSTW data set for the integrated PDFs [26]. The Sivers function at low energy is modeled following what are called the "Bochum" [27] and "Torino" [28] fits in [5].

The model-dependence within Collins' approach manifests itself as dependence on $b_{\max}$ and the fit parameters of, for example, the BLNY model. However within our approach this model-dependence is almost absent. In principle one would need a model from b_c onwards, where the resummed D is no more valid. But the value at which the model-dependence would enter has been shifted to larger values of b by the resummation, such that the remaining model-dependence is negligible. Since the evolution kernel and the TMDs that are evolved are very small in that region, one can set the kernel to zero with no appreciable effect on evolution. Thus no non-perturbative model is needed.

As can be seen in Fig. 2, our implementation of Collins approach is consistent with [4, 5]. From this figure it is clear that if one uses Collins' approach rather than ours, the value of $b_{\max}$ has to be close to 1.5 GeV^{-1} as was indeed found in [15] by a comparison with experimental data.

The definition of quark-TMDs and the new approach to determine the evolution kernel can be extended to gluon-TMDs [29], which can be defined also on-the-light-cone. This new resummation technique can be applied as well to the evolution kernel of the complete hadronic tensor W (built with two TMDs). One major application would be to relate the q_T-spectrum of the Higgs boson production at the LHC at two different energies in a model-independent way.

Acknowledgements: This work is supported by the Spanish MEC, FPA2011-27853-CO2-02. We would like to thank Umberto D'Alesio for useful discussions. The authors have enjoyed discussing our approach for TMDs with V. Braun, M. Diehl, A. Metz and J. Zhou. M.G.E. is supported by the PhD funding program of the Basque Country Government. A.I. and A.S. are supported by BMBF (06RY9191). I.S. is supported by the Ramón y Cajal Program.

---

[*] Electronic address: miguel.gechevarria@fis.ucm.es
[†] Electronic address: ahmad.idilbi@physik.uni-regensburg.de
[‡] Electronic address: andreas.schaefer@physik.uni-regensburg.de
[§] Electronic address: ignazios@fis.ucm.es

[1] D. W. Sivers, Phys. Rev. D **41** (1990) 83.
[2] J. C. Collins, Nucl. Phys. B **396** (1993) 161 [hep-ph/9208213].

[3] V. Barone, F. Bradamante and A. Martin, Prog. Part. Nucl. Phys. **65** (2010) 267 [arXiv:1011.0909 [hep-ph]].

[4] S. M. Aybat and T. C. Rogers, Phys. Rev. D **83** (2011) 114042

[5] S. M. Aybat, J. C. Collins, J. -W. Qiu and T. C. Rogers, Phys. Rev. D **85** (2012) 034043

[6] S. M. Aybat, A. Prokudin and T. C. Rogers,

[7] Z. -B. Kang, B. -W. Xiao and F. Yuan, Phys. Rev. Lett. **107** (2011) 152002

[8] M. Anselmino, M. Boglione and S. Melis, arXiv:1204.1239 [hep-ph].

[9] P. Sun, B. -W. Xiao and F. Yuan, Phys. Rev. D **84** (2011) 094005

[10] B. U. Musch, P. Hägler, M. Engelhardt, J. W. Negele and A. Schäfer, Phys. Rev. D **85** (2012) 094510 [arXiv:1111.4249 [hep-lat]].

[11] M. Garcia-Echevarria, A. Idilbi and I. Scimemi, JHEP **07** (2012) 002

[12] J. Collins, (Cambridge monographs on particle physics, nuclear physics and cosmology. 32)

[13] J. C. Collins and D. E. Soper, Nucl. Phys. B **194** (1982) 445.

[14] J. C. Collins and D. E. Soper, Nucl. Phys. B **193** (1981) 381 [Erratum-ibid. B **213** (1983) 545] [Nucl. Phys. B **213** (1983) 545].

[15] A. V. Konychev and P. M. Nadolsky, Phys. Lett. B **633** (2006) 710 [hep-ph/0506225].

[16] A. V. Manohar and I. W. Stewart, Phys. Rev. D **76** (2007) 074002

[17] P. J. Mulders and R. D. Tangerman, Nucl. Phys. B **461** (1996) 197 [Erratum-ibid. B **484** (1997) 538]

[18] D. Boer and P. J. Mulders, Phys. Rev. D **57** (1998) 5780

[19] A. Idilbi and I. Scimemi, Phys. Lett. B **695** (2011) 463

[20] M. Garcia-Echevarria, A. Idilbi and I. Scimemi, Phys. Rev. D **84** (2011) 011502

[21] S. Moch, J. A. M. Vermaseren and A. Vogt, JHEP **0508** (2005) 049

[22] S. Moch, J. A. M. Vermaseren and A. Vogt, Nucl. Phys. B **688** (2004) 101

[23] J. C. Collins, D. E. Soper and G. F. Sterman, Nucl. Phys. B **250** (1985) 199.

[24] F. Landry, R. Brock, P. M. Nadolsky and C. P. Yuan, Phys. Rev. D **67** (2003) 073016

[25] P. Schweitzer, T. Teckentrup and A. Metz, Phys. Rev. D **81** (2010) 094019 [arXiv:1003.2190 [hep-ph]].

[26] A. D. Martin, W. J. Stirling, R. S. Thorne and G. Watt, Eur. Phys. J. C **63** (2009) 189 [arXiv:0901.0002 [hep-ph]].

[27] J. C. Collins, A. V. Efremov, K. Goeke, S. Menzel, A. Metz and P. Schweitzer, Phys. Rev. D **73** (2006) 014021 [hep-ph/0509076].

[28] M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, S. Melis, F. Murgia, A. Prokudin and C. Turk, Eur. Phys. J. A **39** (2009) 89 [arXiv:0805.2677 [hep-ph]].

[29] P. J. Mulders and J. Rodrigues, Phys. Rev. D **63** (2001) 094021